

NEAR-INFRARED SPECTRAL FEATURES IN SINGLE-AGED STELLAR POPULATIONS

R. P. Schiavon^{1,2}, B. Barbuy³, G. Bruzual A.⁴

ABSTRACT

Synthetic spectra for single-aged stellar populations of metallicities $[M/H] = -0.5, 0.0$ and $+0.5$, ages = 3 to 17 Gyrs, and initial mass function indices $x = 0.1$ to 2.0 were built in the wavelength range $\lambda\lambda$ 6000-10200 Å.

For such we have employed the grid of synthetic spectra described in Schiavon & Barbuy (1999), computed for the stellar parameters $2500 \leq T_{\text{eff}} \leq 6000$ K, $-0.5 \leq \log g \leq 5.0$, $[M/H] = -0.5, 0.0$ and $+0.5$, and $[\alpha/\text{Fe}] = 0.0$, together with the isochrones by Bertelli et al. (1994) and Baraffe et al. (1998).

The behavior of the features NaI λ 8190, CaII λ 8662, TiO λ 6600 and FeH λ 9900 Å in the integrated spectra of single-stellar populations were studied in terms of metallicity, initial mass function and age variations.

The main conclusions are that the NaI doublet is an IMF-sensitive feature, which is however sensitive also to metallicity and age, whereas TiO, CaII and FeH are very sensitive to metallicity and essentially insensitive to IMF and age.

Subject headings: Stars: atmospheres, M giants, globular clusters

1. INTRODUCTION

The analysis of near-infrared (NIR) features in the spectral region $\lambda\lambda$ 6000-10000 Å in the integrated light of globular clusters and galaxies has become of common use to infer

¹Observatório Nacional/CNPq, Departamento de Astronomia, R. Gen. José Cristino 77, 20921-400 Rio de Janeiro, Brazil. Email: ripisc@ucolick.org

²Present address: UCO/Lick Observatory, University of California, Santa Cruz, CA, 95064

³Universidade de São Paulo, IAG, Departamento de Astronomia, C.P. 3386, São Paulo 01060-970, Brazil. Email: barbuy@orion.iagusp.usp.br

⁴Centro de Investigaciones de Astronomía (CIDA), Apartado Postal 264, Mérida 5101-A, Venezuela. Email: bruzual@cida.ve

the parameters of their stellar populations. The NIR spectral region is specially interesting for stellar population studies because M stars (giants and dwarfs) give their maximum contribution to the integrated spectrum of galaxies and clusters in these wavelengths. As an example, NIR line indices which are sensitive to the surface gravity of M stars can be used to constrain the low mass end of the mass function. Also, NIR indices are suitable for deriving metallicities because of their remarkable insensitivity to age. The metallicity calibration of globular clusters using the CaII triplet has been carried out by Bica et al. (1998) for globular clusters in the Galactic bulge, and by Olszewski et al. (1991) for clusters in the Magellanic Clouds.

Long standing problems with the use of IMF-sensitive NIR features in composite systems were discussed in Whitford (1977), Cohen (1978, 1979), Faber & French (1980), Alloin & Bica (1989), Xu et al. (1989), Boroson & Thompson (1991), Delisle & Hardy (1992) and Couture & Hardy (1993), which studied the NaI, CaII and FeH indices in integrated spectra of galaxies and clusters.

In spite of the above mentioned efforts, the strong spectral features which appear in the integrated NIR spectrum of globular clusters and normal galaxies have been far less exploited in stellar population studies than the indices in the optical region, such as Mg₂, Fe5270, and Fe5335 (Burstein et al. 1984; Worthey 1994; Trager et al. 1998). This fact is probably due to two main reasons: the NIR features show a more complex behavior than the ‘green’ features, and the NIR region presents numerous telluric lines which require a proper and careful subtraction.

In this context, spectrum synthesis based on model stellar photospheres and comprehensive line lists can play a key rôle in disentangling the degenerate behaviour of line indices as a function of metallicity, effective temperature and surface gravity. In an attempt to improve our understanding of the NIR features, Schiavon & Barbuy (1999, hereafter SB99) have built an extensive grid of synthetic spectra, based on state-of-the-art model photospheres and molecular and atomic line lists. The NaI ‘doublet’ at $\lambda 8190\text{\AA}$ and the FeH Wing-Ford band (WFB) at $9900\text{\AA}$ (Wing & Ford 1969) were studied as a function of stellar photospheric parameters by Schiavon et al. (1997a, b), whereas the behavior of the TiO bands and the CaII triplet were described in Milone & Barbuy (1994), SB99 and Erderlyi-Mendes & Barbuy (1991).

In this paper we use the grid of high resolution synthetic stellar spectra presented in SB99, to build integrated spectra of single-aged stellar populations (SSPs) in the wavelength range $\lambda\lambda 6000\text{--}10200\text{\AA}$. Previous work with similar aims was presented by Garcia-Vargas et al. (1998) and Milone et al. (1995) where equivalent widths of the CaII triplet and TiO respectively were given.

These models are a useful guide in the interpretation of the integrated spectrum of early-type galaxies, which in the NIR are dominated by M-type stars, and for a better understanding of the age-metallicity degeneracy of old and intermediate age stellar populations.

In §2 we present basic information on the spectrum synthesis of individual stellar models and on the other ingredients required to build the SSP models. In §3 we discuss the behavior of selected NIR features as a function of SSP parameters: metallicity, age and IMF. A summary is presented in §4.

2. INTEGRATED SYNTHETIC SPECTRA OF SINGLE-AGED STELLAR POPULATIONS

2.1. Synthetic Spectra of Individual Stars

SB99 computed a grid of high resolution synthetic spectra in the wavelength range $\lambda\lambda$ 6000 – 10200 Å. The photospheric models by Kurucz (1992), Plez et al. (1992), further unpublished models by Plez (1997), and Allard & Hauschildt (1995) were adopted. The grid covers the following ranges of stellar parameters: $2500 \leq T_{\text{eff}} \leq 6000$ K, $-0.5 \leq \log g \leq 5.0$, $[M/H] = -0.5, 0.0$ and $+0.5$, and $[\alpha/Fe] = 0.0$. The molecular electronic systems included in the calculations are: the CN ($A^2\Pi-X^2\Sigma$) red system, the C₂ ($A^3\Pi_g-X^3\Pi_u$) Swan system, the TiO ($\gamma(A^3\Phi-X^3\Delta)$, $\gamma'(B^3\Pi-X^3\Delta)$, $\delta(b^1\Pi-a^1\Delta)$, $\epsilon(E^3\Pi-X^3\Delta)$, and $\phi(b^1\Pi-d^1\Sigma)$ systems, and the FeH ($A^4\Delta-X^4\Delta$), which includes the WFB. The synthetic spectra were computed in steps of 0.02 Å and rebinned to 1.0 Å with FWHM = 2.0 Å. More details on the atmospheric models and the atomic and molecular data employed are given in SB99.

2.2. Integrated Spectra of Single-aged Stellar Populations

We have combined our grid of synthetic stellar spectra with isochrones collected from the literature to build SSP synthetic spectra. We adopted the Padova isochrones (Bertelli et al. 1994) for $M > 0.6M_{\odot}$, and the isochrones from Baraffe et al. (1998) for $M \leq 0.6M_{\odot}$. We have built integrated spectra of SSPs for $[M/H] = -0.5, 0.0$ and $+0.5$, and two different IMF slopes: $x = 1.35$ (Salpeter) and $x = 2.0$ (dwarf enriched). In Fig. 1 we show examples of the computed spectra in the wavelength range $\lambda\lambda$ 6000-10200 Å, normalized at $\lambda 8157$ Å and convolved with a gaussian of FWHM = 9 Å, for a range of ages (Fig. 1a), metallicities (Fig. 1b) and IMFs (Fig. 1c). As expected, NIR features are stronger in the more metal-rich and older SSPs, since their spectra are dominated by the cooler and redder stars. With a

higher contribution of M dwarfs to the integrated light (Fig. 1c), the spectrum becomes somewhat redder.

In order to test our SSP integrated spectra, we have (1) compared results obtained with Padova and Geneva isochrones, and (2) applied our calculations to the metal-rich globular cluster NGC 6553, as described below.

2.2.1. *Geneva vs. Padova Isochrones*

A check of our results obtained with the Padova isochrones was carried out by using the isochrones from the Geneva group (Schaller et al. 1992 as given in CD-ROM by Leitherer et al. 1996) for stars with $M > 0.6M_{\odot}$, to compute the integrated spectrum of a 13 Gyrs old SSP with solar metallicity and Salpeter IMF. For lower masses we use Baraffe et al.’s isochrones in all cases. In Figures 2 and 3 we compare, respectively, the Geneva and Padova isochrones, and corresponding integrated spectra (normalized at $\lambda = 8170\text{\AA}$).

From Figure 2 it can be seen that the Geneva and Padova isochrones present differences in the red giant branch. While in the first ascent giant branch the Geneva isochrones present a systematically lower temperature ($\sim 100\text{K}$), their red giant tip is much hotter and brighter (for a given temperature) than in the Padova isochrones. The coolest red giant stars in the Geneva isochrone have $T_{\text{eff}} \sim 3300\text{K}$, while in the Padova isochrone red giants can be as cool as $T_{\text{eff}} \sim 2600\text{K}$.

In spite of these large differences in the isochrones, the integrated spectra look remarkably similar in Figure 3. The integrated spectrum computed from the Padova isochrones is brighter than the Geneva one by less than 5% in both ends of the interval. By looking at Figure 11 of SB99, these small differences can be explained by the fact that stars from the tip of the red giant branch (stars later than M4), which are cooler in the Padova isochrones, dominate the integrated light of old SSPs at $\lambda \gtrsim 8500\text{\AA}$, while stars from the warmer part of the red giant branch - RGB (spectral types from mid-K to early-M), which are cooler in the Geneva isochrones, are dominant in the bluer part of the spectral interval under study. The differences in the studied indices are small ($< 10\%$ in equivalent width) except for the Wing-Ford band, for which a 30% difference in equivalent width is found, due to the sensitivity of this feature to the temperature of RGB tip stars.

The fact that we obtain so small spectral differences suggests the existence of a mutual compensation between the above mentioned differences in the isochrones.

2.2.2. The Metal-rich Globular Cluster NGC 6553

Our choice to adopt the Padova set of isochrones is based on our recent study of the T_{eff} scale of M giants in the bulge metal-rich globular cluster NGC 6553 (SB99). The metallicity of NGC 6553 found in the literature ranges from $-0.70 < [\text{Fe}/\text{H}] < +0.47$ (see Table 5 in Barbuy et al. 1999). High resolution spectroscopy tends to give lower values: $-0.7 < [\text{Fe}/\text{H}] < -0.2$, whereas integrated photometry or spectroscopy give higher values: $-0.33 < [\text{Fe}/\text{H}] < +0.47$. Rutledge et al. (1997) gives $[\text{Fe}/\text{H}] = -0.18$ or -0.60 depending whether they use the metallicity scale of globular clusters by Zinn & West (1984) or by Carretta & Gratton (1997). Barbuy et al. (1999) obtained $[\text{Fe}/\text{H}] \approx -0.55$ and $[\alpha/\text{Fe}] \approx +0.4$, which would explain the discrepancy between results obtained from high resolution measurements of Fe lines and integrated spectra where lines of Fe and α -elements give together $[\text{Z}/\text{Z}_{\odot}] \approx 0.0$. Cohen et al. (1999) on the other hand, obtained $[\text{Fe}/\text{H}] \approx -0.2$ and $[\alpha/\text{Fe}] \approx 0.2$. In terms of our present purposes, since the spectra are dominated by the M giants, whose spectra are in turn dominated by TiO bands, and given that both Ti and O are α elements, the use of an overall metallicity of $[\text{M}/\text{H}] \approx 0.0$ is in agreement with the above two results (Barbuy et al. 1999 and Cohen et al. 1999).

The T_{eff} -scale inferred by SB99 for NGC 6553, based on TiO bands, is more compatible with the temperature values in the Padova isochrones, which in particular have a more extended RGB tip; the extension of the RGB of NGC 6553 can be seen in Bruzual et al. (1997) and Guarnieri et al. (1998). Moreover, in a study of the T_{eff} s of M giants from NGC 6528 (another metal rich globular cluster from the Galactic bulge, see Ortolani et al. 1995), we determined $T_{\text{eff}} = 3000$ K for a bonafide M giant of this cluster, which is 300K cooler than the lowest T_{eff} predicted by the Geneva isochrone (Schiavon et al. 1999, in preparation).

3. NaI DOUBLET, CaII TRIPLET, FeH WING-FORD BAND AND TiO BANDS IN SINGLE STELLAR POPULATIONS

In this Section we discuss the effects of metallicity and IMF upon NIR spectral indices commonly used for stellar population diagnosis in integrated spectra of galaxies and clusters (e.g. Cohen 1978, 1979; Delisle & Hardy 1992). The dependence of the TiO bands, NaI doublet, CaII triplet and the WFB on stellar atmospheric parameters have been studied in previous papers (Erdelyi-Mendes & Barbuy 1991; Milone & Barbuy 1994; Schiavon et al. 1997a,b; SB99).

In Table 1 we report the definition of the spectral indices used in this work, including

TiO $\lambda 6600$ Å, NaI $\lambda 8190$ Å, CaII $\lambda 8662$ Å and the FeH WFB at $\lambda 9900$ Å, as given in the literature. In Table 2 we list the equivalent width of the indices measured on the spectra of 13 Gyr SSPs, as a function of metallicity, IMF and age, where convolutions with gaussian profiles of $\text{FWHM} = 9$ and 25 Å were adopted. We note that two definitions of the NaI index are used: Na_{FF} and Na_{S97} as defined in Faber & French (1980, hereafter FF) and Schiavon et al. (1997a, hereafter S97a) respectively.

3.1. CaII Triplet, TiO Bands and FeH WFB

Figures 4a-f show synthetic spectra of 13 Gyr SSPs in the regions of the TiO band ($\lambda 6600$ Å), the WFB ($\lambda 9900$ Å), and CaII ($\lambda 8662$ Å). Figs. 4(a,c,e) correspond to the $[\text{M}/\text{H}] = -0.5, 0.0$ and $+0.5$ models for the Salpeter IMF, and Figs. 4(b,d,f) to the $[\text{M}/\text{H}] = -0.5$ model for the Salpeter and the $x = 2.0$ IMF.

Figs. 4(a,c,e) show that the TiO bands are the most metallicity-sensitive feature under analysis. We restrict our discussion to the TiO_{6600} band, since all bands of the same molecule have essentially the same behavior.

The WFB and the CaII triplet are also very sensitive to metallicity and weakly sensitive to IMF variations. It has been suggested in the literature (Couture & Hardy 1993 and references therein) that the WFB and the CaII triplet (Delisle & Hardy 1992; Jones et al. 1984) are sensitive to the contribution of M dwarfs to the integrated spectrum. Our computations do not confirm this. Fig. 4b shows a residual sensitivity of the TiO_{6600} index to IMF variations, while the WFB (Figs. 4c,d) is only weakly IMF-sensitive, but its response to metallicity is much stronger, being due to the contamination by TiO lines from the (2,3) vibrational band of the δ system.

3.2. The NaI Doublet

Figures 5a,b show spectra of 13 Gyr SSPs in the region of the NaI $\lambda 8190$ Å feature. Fig. 5a correspond to the $[\text{M}/\text{H}] = -0.5, 0.0$ and $+0.5$ models for the Salpeter IMF, and Fig. 5b to the $[\text{M}/\text{H}] = -0.5$ model for the Salpeter and the $x = 2.0$ IMF.

Figs. 5a,b show that the NaI doublet is by far the most IMF-sensitive index studied here. The Na_{S97} index displays a strong dependence on IMF (Table 2). It also shows a dependence on metallicity. Besides, the NaI doublet is contaminated by TiO lines (which are very sensitive to metallicity) at the red side of the feature.

It has been suggested in the literature (Cohen 1978; Xu et al. 1989; Alloin & Bica 1989; Terndrup et al. 1990; Delisle & Hardy 1992) that the NaI doublet is more sensitive to metallicity than to IMF variations. Figs. 5 a,c show that the NaI *lines* are indeed strongly sensitive to metallicity. However, the NaI *index*, as defined in the literature (Table 1), has its metallicity-sensitivity reduced by the TiO lines that contaminate the red continuum window, which is thus lowered in the spectra of metal-rich stellar populations.

Figs. 5 and the values in Table 2 suggest that the NaI index is very sensitive to IMF. In Fig. 6 the equivalent width of NaI, according to the definitions by FF and Schiavon et al. (1997) and CaII lines (normalized by the values corresponding to a SSP of 13Gyrs, solar metallicity and Salpeter IMF), are plotted against the exponent of a power-law IMF, for different metallicities and an age of 13Gyrs. The NaI index, in both definitions, is as sensitive to IMF as to metallicity. We therefore conclude that the possibility that an enhancement of the NaI lines may be due to a dwarf-enriched IMF cannot be excluded. An uncertainty in metallicity of $\Delta[M/H] = 0.25$ dex translates into an uncertainty of 15% in x for $x > 1$, and 50% for $x < 1$; therefore the use of the NaI doublet as an IMF indicator requires narrow constraints on the average metallicity of the stellar population.

Figs. 5 suggest that in the integrated spectra of galaxies, where lines are broadened by stellar velocity dispersion, IMF and metallicity effects may be distinguished by their different influence on the line shape, as proposed by Boroson & Thompson (1991). A higher metallicity leads to a profile with a deeper red side, while a profile with a deeper blue side can be due to either a stronger metallicity or a dwarf-enriched IMF.

3.3. Sensitivity of the Features to Age

In Table 3 are given computed synthetic indices convolved with $\text{FWHM} = 25\text{\AA}$ for single stellar populations for ages $3 < \text{Age (Gyr)} < 17$ and $-0.5 < [M/H] < +0.5$ for a Salpeter IMF.

In Fig. 7 we plot the four indices versus the age of SSPs for $[M/H] = -0.5$ and 0.0 . It appears that the TiO band, the CaII triplet, and the WFB are essentially insensitive to age. Note the strong metallicity dependence of the TiO bands. The NaI doublet is very sensitive to age, as well as to the IMF slope and metallicity, which makes the behavior of this feature quite complex. It is of interest to explore indices such as TiO bands and the CaII triplet to help disentangle the age-metallicity degeneracy in composite stellar populations.

4. SUMMARY

We have built high resolution synthetic spectra of SSPs in the wavelength range $\lambda\lambda$ 6000-10200 Å for SSPs of $[M/H] = -0.5, 0.0$ and $+0.5$, and IMF index $x = 0.1-2.0$. The basic ingredients for such SSP models are the isochrones by Bertelli et al. (1994) and Baraffe et al. (1998), combined to a grid of synthetic spectra computed for a wide range of stellar parameters (SB99).

In order to make our computations useful for observers, we present a table of TiO $\lambda 6600$ Å, NaI $\lambda 8190$ Å, CaII $\lambda 8662$ Å and the FeH WFB at $\lambda 9900$ Å indices in single stellar populations, as a function of metallicity, IMF and age.

The main conclusions of this paper are:

1) We have shown that the TiO bands, the CaII triplet, and the WFB are very sensitive to metallicity and essentially invariant with IMF and age. 2) The NaI doublet is far more intense for a dwarf-enriched than for the Salpeter IMF, and the feature is also affected by metallicity and age variations. 3) The better understanding of the NIR indices as a function of metallicity, age and IMF obtained through the present spectrum synthesis computations, provides us with tools, which combined to optical indices, can help disentangle the age - metallicity - IMF degeneracy in composite stellar systems.

We are grateful to the referee for helpful comments on an earlier version of this paper. Calculations were carried out in a DEC-Alpha 3000/700 workstation. RPS acknowledges support from CNPq fellowship No. 3000098/4.

REFERENCES

- Allard, F., Hauschildt, P.H. 1995, ApJ, 445, 433
Alloin, D., Bica, E. 1989, A&A, 217, 57
Baraffe, I., Chabrier, G., Allard, F., Hauschildt, P.H., 1998, A&A, 337, 403
Barbuy, B., Renzini, A., Ortolani, S., Bica, E., Guarnieri, M.D. 1999, 341, 539
Bertelli, G., Bressan, A., Chiosi, C., Fagotto, F., Nasi, E. 1994, A&AS, 106, 275
Bica, E., Clariá, J.J., Piatti, A.E., Bonatto, C. 1998, A&AS, 131, 483
Borson, T.A., Thompson, I.B. 1991, AJ, 101, 111
Bruzual, G., Barbuy, B., Ortolani, S., Bica, E., Cuisinier, F., Lejeune, T., Schiavon, R. 1997, AJ, 114, 1531

- Burstein, D., Faber, S.M., Gaskell, C.M., Krumm, N. 1984, ApJ, 287, 586
- Carretta, E., Gratton, R.G. 1997, A&AS, 121, 95
- Cohen, J.G. 1978, ApJ, 221, 788
- Cohen, J.G. 1979, ApJ, 228, 405
- Cohen, J.G., Gratton, R.G., Behr, B.B., Carretta, E. 1999, ApJ, in press.
- Couture, J., Hardy, E. 1993, ApJ, 406, 142
- Delisle, S., Hardy, E. 1992, AJ, 103, 711
- Erdelyi-Mendes, M., Barbuy, B. 1991, A&A, 241, 176
- Faber, S.M., French, H. 1980, ApJ, 235, 405 (FF)
- Garcia-Vargas, M.L., Molla, M., Bressan, A. 1998, A&AS, 130, 513
- Guarnieri, M.D., Ortolani, S., Montegriffo, P., Renzini, A., Barbuy, B., Bica, E., Moneti, A. 1998, A&A, 331, 70
- Jones, J.E., Alloin, D.M., Jones, B.J.T., 1984, ApJ, 283, 457.
- Kurucz, R. 1992, in IAU Symp. 149, The Stellar Populations of Galaxies, eds. B. Barbuy & A. Renzini, Kluwer Acad. Pub., p. 225
- Leitherer, C. et al. 1996, PASP, 108, 996
- Milone, A., Barbuy, B. 1994, A&AS, 108, 449
- Milone, A., Barbuy, B., Bica, E. 1995, A&AS, 113, 547
- Olszewski, E.W., Schommer, R.A., Suntzeff, N.B., Harris, H.C. 1991, AJ, 101, 515
- Ortolani, S., Renzini, A., Gilmozzi, R., Marconi, G., Barbuy, B., Bica, E., Rich, R.M. 1995, Nature, 377, 701.
- Plez, B. 1997, private communication
- Plez, B., Brett, J.M., Nordlund, Å. 1992, A&A, 256, 551
- Rutledge, G.A., Hesser, J.E., Stetson, P.B. 1997, PASP, 109, 907
- Schaller, G., Schaerer, D., Meynet, G., Maeder, A. 1992, A&AS, 96, 269
- Schiavon, R.P., Barbuy, B., Rossi, S., Milone, A. 1997a, ApJ, 479, 902 (S97)
- Schiavon, R.P., Barbuy, B., Singh, P.D. 1997b, ApJ, 484, 499.
- Schiavon, R.P., Barbuy, B. 1999, ApJ, 510, 934 (SB99)
- Schiavon, R.P., Barbuy, B., Bica, E. 1999, in preparation
- Terndrup, D.M., Frogel, J.A., Whitford, A.E. 1990, ApJ, 357, 453

- Trager, S.C., Worthey, G., Faber, S.M., Burstein, D., González, J.J. 1998, ApJS, 116, 1
- Xu, Z., Véron-Cetty, M.-P., Véron, P. 1989, A&A, 211, L12
- Whitford, A.E., 1977, ApJ, 211, 527.
- Wing, R.F., Ford, W.K., Jr. , 1969, PASP, 81, 527.
- Worthey, G., 1994, ApJS, 95, 107.
- Zinn, R., West, M.J., 1984, ApJS, 55, 45.

Table 1. Definition of spectral indices measured

Index	Blue continuum	Bandpass	Red continuum
EW_{6600}	6512.1-6538.1	6617.2-6992.5	7036.9-7048.0
EW_{8190}^{FF}	8169.0-8171.0	8172.0-8209.0	8209.0-8211.0
EW_{8190}^{S97}	8171.5-8172	8172-8197	8233.5-8234.2
EW_{Ca8662}	8637.2-8646.2	8653.2-8668.4	8847.6-8854.0
EW_{WFB}	9891.8-9895.1	9895.1-9958.6	9958.6-9962.2

Table 2. Spectral indices (equivalent widths in Å) in synthetic Single Stellar Populations of 13 Gyr, for convolutions of 9 and 25 Å

[M/H]	x	Na _{FF}	Na _{S97}	TiO ₆₆₀₀	Ca ₈₆₆₂	WFB
FWHM=9Å						
−0.5	1.35	1.03	0.95	9.3	1.41	0.56
0.0	1.35	1.15	1.19	24.9	1.57	1.35
+0.5	1.35	1.45	1.48	34.0	1.80	2.03
−0.5	2.0	1.56	1.40	11.2	1.41	0.72
0.0	2.0	1.78	1.70	28.5	1.50	1.52
FWHM=25Å						
−0.5	1.35	0.47	0.55	7.48	0.52	0.28
0.0	1.35	0.51	0.69	21.7	0.78	0.72
+0.5	1.35	0.67	0.88	29.7	0.90	1.03
−0.5	2.0	0.80	0.81	9.38	0.60	0.45
0.0	2.0	0.90	0.99	25.3	0.69	0.90

Table 3. Spectral indices (equivalent widths in Å in synthetic Single Stellar Populations as a function of age and metallicity, for a Salpeter IMF, adopting a convolution of FWHM = 25

age	[M/H]	Na _{FF}	Na _{S97}	TiO ₆₆₀₀	Ca ₈₆₆₂	WFB
3	-0.5	0.532	0.578	0.379	0.749	0.339
5	-0.5	0.634	0.633	0.377	0.644	0.381
6	-0.5	0.676	0.663	0.387	0.689	0.393
8	-0.5	0.723	0.691	0.397	0.691	0.391
10	-0.5	0.799	0.736	0.385	0.599	0.411
12	-0.5	0.886	0.773	0.326	0.717	0.379
13	-0.5	0.925	0.796	0.345	0.663	0.394
15	-0.5	0.992	0.832	0.341	0.748	0.407
17	-0.5	1.073	0.878	0.319	0.640	0.401
3	0.0	0.422	0.660	0.809	0.926	0.884
5	0.0	0.558	0.739	0.864	0.993	0.886
6	0.0	0.647	0.788	0.893	1.037	0.961
8	0.0	0.698	0.828	0.928	0.955	0.975
10	0.0	0.784	0.879	0.971	0.946	1.020
12	0.0	0.888	0.940	0.970	0.908	1.019
13	0.0	1.000	1.000	1.000	1.000	1.000
15	0.0	1.077	1.034	0.945	0.891	0.957
17	0.0	1.071	1.043	1.058	1.016	1.010
3	0.5	0.530	0.813	0.892	1.178	1.005
5	0.5	0.673	0.900	0.994	1.174	1.163
6	0.5	0.775	0.966	1.128	1.193	1.264
8	0.5	0.912	1.038	1.139	1.276	1.246
10	0.5	0.973	1.078	1.237	1.080	1.280
12	0.5	1.220	1.214	1.303	0.895	1.364
13	0.5	1.315	1.272	1.367	1.150	1.436
15	0.5	1.353	1.299	1.467	1.164	1.514
17	0.5	1.406	1.325	1.403	1.051	1.353

FIGURE CAPTIONS

Fig. 1.— Synthetic spectra in the wavelength range $\lambda\lambda$ 6000 - 10200 Å for (a) Salpeter IMF, $[M/H] = 0.0$ and ages of 3 Gyr (solid line), 8 Gyr (dashed line) and 13 Gyr (dotted line); (b) Salpeter IMF and $[M/H] = -0.5$ (solid line), 0.0 (dashed line) and +0.5 (dotted line), normalized at $\lambda 8170$ Å; (c) $[M/H]$ and IMF index $x = 1.35$ (solid line) and 2.0 (dashed line).

Fig. 2.— Isochrones of 13Gyr and $[Z/Z_{\odot}] = 0.0$ from the Padova (open circles) and Geneva groups (filled squares).

Fig. 3.— Integrated spectra generated with the use of Geneva (solid line) and Padova (dotted line) isochrones (normalized at $\lambda = 8170$ Å) are compared.

Fig. 4.— Synthetic spectra for single stellar populations with 13Gyr, a Salpeter IMF and $[M/H] = -0.5$ (solid), 0.0 (dashed) and +0.5 (dotted) for the features (a) TiO_{6600} , (c) WFB, (e) CaII triplet; and synthetic spectra for SSPs of $[M/H] = -0.5$ for Salpeter (solid) and dwarf-enriched (dashed) IMFs for (b) the TiO_{6600} , (d) WFB, (f) CaII triplet.

Fig. 5.— Synthetic spectra of the NaI feature for SSPs of 13 Gyr, Bertelli et al. (1994) isochrones, and (a) Salpeter IMF and $[M/H] = -0.5$ (solid), 0.0 (dashed) and +0.5 (dotted); (b) $[M/H] = -0.5$ and IMF index $x = 1.35$ (solid) and 2.0 (dashed).

Fig. 6.— CaII, NaI_{S97} and NaI_{FF} vs. IMF index x , for $[M/H] = -0.5$, 0.0 and +0.5 and age=13Gyrs.

Fig. 7.— Indices for TiO_{6600} , WFB, NaI and CaII measured on the spectra of single stellar populations as a function of age (Gyrs) for $[M/H] = -0.5$ (solid lines) and 0.0 (dotted lines). Equivalent widths are normalized to the value corresponding to a SSP with solar metallicity 13Gyrs and a Salpeter IMF.













